

VI - CLAIMING

What is claimed is :

- 1 – A quantitative method (quantifying by means of Biolumager or ELISA with colorimetric detection) for the measurement of specific mRNA utilizing the labeled nucleotide probes 1 and 2 (labeling with ^{32}P -dCTP and with biotin) directed at exons 7 and 8 respectively of the SMN gene for the molecular diagnosis of SMA.
- 2 – A quantitative method according to claim 1 may be used in the control of mRNA concentrations in the future gene therapy of patients with SMA.
- 3 – A quantitative method based on the measurement of specific mRNA for the construction of molecular diagnostic kits and for the control of mRNA concentrations in the future gene therapy for genetic disorders associated to a deletion or mutation of the gene(s) such as Duchene myopathies, mucoviscidose, or to a duplication of the gene such as Charcot-Marie-Tooth disease type 1A.

VII - REFERENCES

- 1) Sambrook J, Fritsch EF, Maniatis T. Extraction, purification and analysis of messenger RNA from eukariotic cells. In : Molecular cloning, a laboratory manual, 2nd ed. Cold Spring Harbor, NY : Cold Spring Harbor Laboratory Press, 1989 : 7.28 - 7.52.
- 2) Petersen NE, Larsen LK, Nissen H, Jensen A, Petersen H *et al.* Improved RNase protection assay for quantifying LDL receptor mRNA ; estimation of analytical imprecision and biological variance in peripheral blood mononuclear cells. *Clin Chem* 1995 ; **41** : 1605 - 1613.
- 3) Saiki RK, Scharf S, Faloona F, Mullis KB, Horn CT, Erlich HA, Arnheim N. Enzymatic amplification of β -globine genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 1985 ; **230** : 1350 - 11354.
- 4) Kawasaki ES, Wang AM. Detection of gene expression. In : PCR technology, Erlich H.A., ed., New York : Stockton, 1989 : 89 - 97..
- 5) Radnsky R, Bucana CD, Ellis LM, Sanchez R, Cleary KR, Brigati DJ, Fidler IJ. A rapid colorimetric *in situ* messenger RNA hybridization technique for analysis of epidermal growth factor receptor in paraffin-embedded surgical specimens of human colon carcinomas. *Cancer Res* 1993 ; **53** : 937 - 943.
- 6) Murray GI. *In situ* PCR. *J Pathol* 1993 ; **169** : 187 - 188.
- 7) Fox JL, Hsu PH, Legator MS, Morrison LE, Seelig SA. Fluorescence *in situ* hybridization : powerful molecular tool for cancer prognosis. *Clin Chem* 1995 ; **41** : 1554 - 1559.
- 8) Stanley PE. A survey of more than 90 commercially available luminometers and imaging devices for low-light measurements of chemiluminescence and bioluminescence, including instruments for manual, automatic and specialized operation, for HPLC, LC, GLC and microtitre plates. Part 1 : descriptions. *J Biolumin Chemilumin* 1992 ; **7** : 77 - 108.

- 9) Van Doornum GJ, Buimer M, Prius M, Henquet CJ, Coutinho RA, Plier PK *et al.*
Detection of Chlamydia trachomatis infection in urine samples from men and
women by ligase chain reaction. *J Clin Microbiol* 1995 ; **33** : 2042 - 2047.
- 10) Duck P, Alvarado-Urbina G, Burdick B, Collier B. Probe amplifier system based on
chimeric cycling oligonucleotides. *Biotechniques* 1990 ; **9** : 142 - 148.
- 11) Walker GT, Little MC, Nadeau JG, Shank DD. Isothermal in vitro amplification of
DNA by a restriction enzyme/DNA polymerase system. *Proc Natl Acad Sci USA*
1992 ; **89** : 392 - 396.
- 12) Pearn J. The gene frequency of acute Werdnig-Hoffmann disease (SMA type I). A
total population survey in North East England. *J Med Genet* 1973 ; **10** : 260 - 265.
- 13) Pearn J. Incidence, prevalence, and gene frequency studies of chronic childhood
spinal muscular atrophy. *J Med Genet* 1978 ; **15** : 409 - 413.
- 14) Czeizel A, Hamula J. A Hungarian study on Werdnig-Hoffmann disease. *J Med*
Genet 1989 ; **26** : 761 - 763.
- 15) Munsat TL, Workshop report. International SMA collaboration. *Neuromusc Disord*
1991 ; **1** : 81.
- 16) Lefebvre S, Bürglen L, Reboullet S *et al.* Identification and characterization of a
spinal muscular atrophy-determining gene. *Cell* 1995 ; **80** : 155 - 165.
- 17) Van der Steege G, Grootsholten PM, Van der Vlies *et al.* PCR-based DNA test to
confirm clinical diagnosis of autosomal recessive spinal muscular atrophy. *Lancet*
1995 ; **345** : 985 - 986.
- 18) Askanas V, Engel WK. New program for investigating adult human skeletal muscle
growth aneurally in tissue culture. *Neurology* 1975 ; **25** : 58 - 67.
- 19) Askanas V, Gallez-Hawkins EF. Synergistic influence of polypeptide growth factors
on cultured human muscle. *Arch Neurol* 1985 ; **42** : 749 - 752.
- 20) Ann DK, Wu MM, Huang T, Carlson DM, Wu R. Retinol-regulated gene expression
in human tracheobronchia epithelial cells. Enhanced expression of elongation

factor EF-1alpha. *J Biol Chem* 1988 ; **263** : 3546 - 3549.

- 21) Southern EM. Detection of specific sequences among DNA fragments separated by gel electrophoresis. *Mol Biol* 1975 ; **98** : 503 - 517.